## AMENDMENTS TO THE CLAIMS

## 1. (Canceled)

- 2. (Original) A crosslinking agent or a curing agent for resins, the agent containing as an active component a polyacrylic hydrazide having an average molecular weight of 70,000 to 150,000, a hydrazide conversion ratio of at least 45% and 400 or more hydrazide groups in one molecule.
- **3.** (Original) A crosslinking agent or a curing agent for resins, the agent containing as an active component a polyacrylic hydrazide having an average molecular weight of 80,000 to 110,000, a hydrazide conversion ratio of at least 45% and 450 or more hydrazide groups in one molecule.
- **4.** (Original) A crosslinking agent or a curing agent for resins, the agent containing as an active component a polyacrylic hydrazide having an average molecular weight of 80,000 to 90,000, a hydrazide conversion ratio of at least 50% and 500 or more hydrazide groups in one molecule.
- **5.** (Original) A crosslinking agent or a curing agent for resins, the agent containing as an active component a polyacrylic hydrazide having an average molecular weight of 20,000 to 40,000, a hydrazide conversion ratio of at least 65% and 150 or more hydrazide groups in one molecule.
- 6. (Original) A crosslinking agent or a curing agent for resins, the agent containing as an active component a polyacrylic hydrazide having an average molecular weight of 20,000 to 35,000, a hydrazide conversion ratio of at least 65% and 150 or more hydrazide groups in one molecule.
- 7. (Currently Amended) A resin composition comprising at least one kind of resin selected from an acrylic resin having at least one carbonyl group in the molecule, a

urethane resin and an epoxy resin, and the crosslinking agent or the curing agent as defined in claims [1] 2 to 6.

- **8.** (Currently Amended) A crosslinked or cured product formed by crosslinking or curing at least one kind of resin selected from an acrylic resin having at least one carbonyl group in the molecule, a urethane resin and an epoxy resin using the crosslinking agent or the curing agent as defined in claims [[1]] 2 to 6.
- **9. (Original)** A polyacrylic hydrazide having an average molecular weight of 20,000 to 30,000, and a hydrazide conversion ratio of at least 70%.
- **10.** (Currently Amended) A polyacrylic hydrazide having an average molecular weight of 50,000 to 150,000 70,000 to 86,000, and a hydrazide conversion ratio of at least 50%.